

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 April 2004 (15.04.2004)

PCT

(10) International Publication Number
WO 2004/031181 A1

(51) International Patent Classification⁷: C07D 409/12,
413/14, 417/14, 409/14, A61K 31/55, A61P 25/00

GlaxoSmithKline, New Frontiers Science Park South,
Third Avenue, Harlow, Essex CM19 5AW (GB).

(21) International Application Number:
PCT/EP2003/011174

(74) Agent: MCKINNELL, Denise; GlaxoSmithKline, CIP
(CN925.1), 980 Great West Road, Brentford, Middlesex
TW8 9GS (GB).

(22) International Filing Date: 6 October 2003 (06.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0223226.2	7 October 2002 (07.10.2002)	GB
0223236.1	7 October 2002 (07.10.2002)	GB
0314627.1	23 June 2003 (23.06.2003)	GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): FORBES, Ian, Thomson [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). GRIBBLE, Andrew, Derrick [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). LIGHT-FOOT, Andrew, P. [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). PAYNE, Andrew, H. [GB/GB];

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: SULFONAMIDE DERIVATIVES AS ANTIPSYCHOTIC AGENTS



(I)


(57) Abstract: The invention provides compounds of formula (I) wherein A and B represent the groups $-(CH_2)_m-$ and $-(CH_2)_n-$ respectively; R^1 represents C_{1-6} alkyl; R^2 represents hydrogen, halogen, hydroxy, cyano, nitro, hydroxy C_{1-6} alkyl, trifluoromethyl, trifluoromethoxy, C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_pC_{3-6}$ cycloalkyl, $-(CH_2)_pOC_{3-6}cy-$

cloalkyl, $-COC_{1-6}alkyl$, $-SO_2C_{1-6}alkyl$, $-SOC_{1-6}alkyl$, $-S-C_{1-6}alkyl$, $-CO_2C_{1-6}alkyl$, $-CO_2NR^4R^5$, $-SO_2NR^4R^5$, $-(CH_2)_pNR^4R^5$, $-(CH_2)_pNR^4COR^5$, an optionally substituted aryl group, an optionally substituted heteroaryl group or an optionally substituted heterocyclyl group; R^3 represents hydrogen or C_{1-6} alkyl; Ar^1 represents an optionally substituted heteroaryl group; Ar^2 represents an optionally substituted phenyl or an optionally substituted heteroaryl group; Y represents a bond, $-O-$, $-C_{1-6}alkyl-$, $-CR^6R^7X-$, $-XCR^6R^7-$, $-NR^8CO-$ or $-CONR^8-$; X represents oxygen, sulfur, $-SO-$ or $-SO_2-$; R^4 and R^5 each independently represent hydrogen or $C_{1-6}alkyl$ or, together with the nitrogen or other atoms to which they are attached, form an azacycloalkyl ring or an oxo-substituted azacycloalkyl ring; R^6 and R^7 each independently represent hydrogen, $C_{1-6}alkyl$ or fluoro; R^8 represents hydrogen or $C_{1-6}alkyl$; m and n independently represent an integer selected from 1 and 2; p independently represents an integer selected from 0, 1, 2 and 3; or a pharmaceutically acceptable salt, solvate or pharmaceutically acceptable derivative thereof. The compounds are useful in therapy, in particular as antipsychotic agents.

SULFONAMIDE DERIVATIVES AS ANTIPSYCHOTIC AGENTS


This invention relates to novel compounds, pharmaceutical compositions containing them and their use in therapy, in particular as antipsychotic agents.

US patent number US 5,684,195 discloses a method of preparing sulfonamides of the formula



wherein R^{10} represents, *inter alia*, an optionally substituted 5- or 6-membered heteroaryl, R^{20} represents, *inter alia*, tetrahydroisoquinoliny and R^{30} represents, *inter alia*, hydrogen or C_{1-6} alkyl. In US 5,684,195, it is stated that sulfonamides in general have been widely used for the treatment of bacterial or viral infections and are also found in drugs such as diuretics, hypoglycemic and antimalarial agents amongst others.

According to the invention, there is provided a compound of formula (I):



wherein

A and B represent the groups $-(CH_2)_m-$ and $-(CH_2)_n-$ respectively;

R^1 represents C_{1-6} alkyl;

R^2 represents hydrogen, halogen, hydroxy, cyano, nitro, hydroxy C_{1-6} alkyl, trifluoromethyl, trifluoromethoxy, C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_pC_{3-6}$ cycloalkyl, $-(CH_2)_pOC_{3-6}$ cycloalkyl, $-COC_{1-6}$ alkyl, $-SO_2C_{1-6}$ alkyl, $-SOC_{1-6}$ alkyl, $-SC_{1-6}$ alkyl, $-CO_2C_{1-6}$ alkyl, $-CO_2NR^4R^5$, $-SO_2NR^4R^5$, $-(CH_2)_pNR^4R^5$, $-(CH_2)_pNR^4COR^5$, an optionally substituted aryl group, an optionally substituted heteroaryl group or an optionally substituted heterocyclyl group;

R^3 represents hydrogen or C_{1-6} alkyl;

Ar^1 represents an optionally substituted heteroaryl group;

Ar^2 represents an optionally substituted phenyl or an optionally substituted heteroaryl group;

Y represents a bond, $-O-$, $-C_{1-6}$ alkyl-, $-CR^6R^7X-$, $-XCR^6R^7-$, $-NR^8CO-$ or $-CONR^8-$;

X represents oxygen, sulfur, $-SO-$ or $-SO_2-$;

R^4 and R^5 each independently represent hydrogen or C_{1-6} alkyl or, together with the nitrogen or other atoms to which they are attached, form an azacycloalkyl ring or an oxo-substituted azacycloalkyl ring;

R^6 and R^7 each independently represent hydrogen, C_{1-6} alkyl or fluoro;

R⁸ represents hydrogen or C₁₋₆alkyl;

m and n independently represent an integer selected from 1 and 2;

p independently represents an integer selected from 0, 1, 2 and 3;

5 or a pharmaceutically acceptable salt, solvate or pharmaceutically acceptable derivative thereof.

10 In a first aspect of the invention, there is provided a compound of the formula (I) as defined above with the proviso that when m represents 1 and n represents 2 or m represents 2 and n represents 1 and R² represents halogen, C₁₋₆alkyl or C₁₋₆alkoxy, Y is other than a bond.

It is to be understood that the present invention covers all combinations of particular and preferred groups described herein above.

15 As used herein, the term "alkyl", either alone or as part of another group, refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms. For example, C₁₋₆alkyl means a straight or branched alkyl containing at least 1, and at most 6, carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, isobutyl, isopropyl, t-butyl and 1,1-dimethylpropyl.

20 As used herein, the term "alkoxy" refers to a straight or branched alkoxy group containing the specified number of carbon atoms. For example, C₁₋₆alkoxy means a straight or branched alkoxy group containing at least 1, and at most 6, carbon atoms. Examples of "alkoxy" as used herein include, but are not limited to, methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy or hexyloxy.

30 As used herein, the term "cycloalkyl" refers to a non-aromatic hydrocarbon ring containing the specified number of carbon atoms. For example, C₃₋₇cycloalkyl means a non-aromatic ring containing at least three, and at most seven, ring carbon atoms. Examples of "cycloalkyl" as used herein include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. A C₆₋₇cycloalkyl group is preferred.

35 As used herein, the term "halogen" refers to the elements fluorine, chlorine, bromine and iodine. Preferred halogens are fluorine, chlorine and bromine.

40 As used herein, the term "aryl" refers to a phenyl or a naphthyl ring.

As used herein, the term "heteroaryl" refers to a 5- or 6-membered heterocyclic aromatic ring or a fused bicyclic heteroaromatic ring system.

As used herein, the term "heterocyclyl" refers to a 3- to 7-membered monocyclic saturated ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Examples of suitable heterocyclic rings include, but are not limited to, piperidine and morpholine.

5

As used herein, the term "5- or 6-membered heterocyclic aromatic ring" refers to a monocyclic unsaturated ring containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Examples of suitable 5- and 6-membered heterocyclic aromatic rings include, but are not limited to, furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl, pyrazolyl, isothiazolyl and isoxazolyl.

10

As used herein, the term "fused bicyclic heteroaromatic ring system" refers to a ring system comprising one six-membered unsaturated ring and one 5- or 6-membered unsaturated ring fused together, the ring system containing at least one heteroatom independently selected from oxygen, nitrogen and sulfur. Examples of suitable fused bicyclic heteroaromatic ring systems include, but are not limited to, indolyl, indolynyl, benzofuranyl, benzothienyl, quinolyl, isoquinolyl, tetrahydroquinolyl, benzodioxanyl, indanyl and tetrahydronaphthyl.

15

20

As used herein, the term "azacycloalkyl ring" refers to a 4- to 7-membered monocyclic saturated ring containing one nitrogen atom. Examples of suitable azacycloalkyl rings are azetidine, pyrrolidine, piperidine and azepine.

25

As used herein, the term "oxo-substituted azacycloalkyl ring" refers to an azacycloalkyl ring as defined above substituted by one oxo group. Examples of suitable oxo-substituted azacycloalkyl rings include, but are not limited to, azetidinone, pyrrolidinone, piperidinone and azepinone.

30

As used herein, the term "optionally substituted" refers to optional substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

35

Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer or step or group of integers but not to the exclusion of any other integer or step or group of integers or steps.

40

As used herein "pharmaceutically acceptable derivative" includes any pharmaceutically acceptable salt, solvate, ester or salt of such ester of a compound of formula (I) which, upon administration to the recipient is capable of providing

(directly or indirectly) a compound of formula (I) or an active metabolite or residue thereof.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula (I) or a salt thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include water, methanol, ethanol and acetic acid. Most preferably the solvent used is water and the solvate may also be referred to as a hydrate.

It will be appreciated that for use in medicine the salts of formula (I) should be pharmaceutically acceptable. Suitable pharmaceutically acceptable salts will be apparent to those skilled in the art and include for example acid addition salts formed with inorganic acids e.g. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and organic acids e.g. succinic, maleic, malic, mandelic, acetic, fumaric, glutamic, lactic, citric, tartaric, benzoic, benzenesulfonic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid. Other non-pharmaceutically acceptable salts e.g. oxalates, may be used, for example in the isolation of compounds of formula (I) and are included within the scope of this invention. Also included within the scope of the invention are solvates and hydrates of the compounds of formula (I).

Certain of the compounds of formula (I) may form acid addition salts with one or more equivalents of the acid. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms thereof.

Certain compounds of formula (I) may exist in stereoisomeric forms (e.g. they may contain one or more asymmetric carbon atoms). The individual stereoisomers (enantiomers and diastereomers) and mixtures of these are included within the scope of the present invention. The present invention also covers the individual isomers of the compounds represented by formula (I) as mixtures with isomers thereof in which one or more chiral centres are inverted. Likewise, it is understood that compounds of formula (I) may exist in tautomeric forms other than that shown in the formula and these are also included within the scope of the present invention.

Preferably, R^1 represents C_{1-4} alkyl. More preferably, R^1 represents methyl, ethyl, n-propyl or isopropyl. Even more preferably, R^1 represents methyl.

The group R^2 may be located at any free position on its respective phenyl ring.

When R^2 represents an optionally substituted aryl group, an optionally substituted heteroaryl group or an optionally substituted heterocyclyl group, the optional substituents may be independently selected from C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_p NR^4 R^5$ wherein p, R^4 and R^5 are as hereinbefore defined, halogen, trifluoromethyl,

trifluoromethoxy, cyano and $-S-C_{1-6}alkyl$. Preferably, the optional substituents are independently selected from chloro, fluoro, bromo, methyl, ethyl, t-butyl, methoxy, trifluoromethyl, trifluoromethoxy, cyano and $-S-methyl$.

- 5 Preferably, R^2 represents hydrogen, halogen, $C_{1-6}alkyl$, $C_{1-6}alkoxy$ or $-(CH_2)_pNR^4R^5$ wherein p , R^4 and R^5 are as hereinbefore defined. More preferably, R^2 represents hydrogen, halogen, $C_{1-4}alkyl$, $C_{1-4}alkoxy$ or dimethylamino. Even more preferably, R^2 represents hydrogen or methoxy.
- 10 Preferably, R^3 represents hydrogen or $C_{1-4}alkyl$. More preferably, R^3 represents hydrogen, methyl, ethyl, n-propyl or isopropyl. Even more preferably, R^3 represents hydrogen.

- 15 When Ar^1 represents an optionally substituted heteroaryl group, the optional substituents may be independently selected from $C_{1-6}alkyl$, $C_{1-6}alkoxy$, halogen, trifluoromethyl, trifluoromethoxy, cyano and $-S-C_{1-6}alkyl$. Preferably, the optional substituents are independently selected from chloro, fluoro, methyl, methoxy, trifluoromethyl and trifluoromethoxy.

- 20 Preferably, Ar^1 is substituted by 0 to 3 substituents, more preferably 0, 1 or 2 substituents.

Preferably, Ar^1 represents optionally substituted thienyl.

- 25 Preferably, Ar^1 represents optionally substituted thienyl whereby one or more optional substituents are selected from halogen (such as chloro, e.g. 4-chloro, or fluoro, e.g. 4-fluoro, 2,4-difluoro or 3,4-difluoro), $C_{1-6}alkyl$ (such as methyl, e.g. 2-methyl), $C_{1-6}alkoxy$ (such as methoxy, e.g. 3-methoxy or 4-methoxy), trifluoromethyl (e.g. 3-trifluoromethyl or 4-trifluoromethyl) and trifluoromethoxy. Other examples of
- 30 multiple optional substituents include, for example, 2-methyl-4-chloro. Even more preferably, Ar^1 represents unsubstituted thienyl.

- When Ar^2 represents optionally substituted phenyl or an optionally substituted heteroaryl group, the optional substituents may be independently selected from $C_{1-6}alkyl$, $C_{1-6}alkoxy$, halogen, trifluoromethyl, trifluoromethoxy, cyano and $-S-C_{1-6}alkyl$.
- 35 Preferably, the optional substituents are independently selected from chloro, fluoro, methyl, methoxy, trifluoromethyl and trifluoromethoxy.

- 40 Preferably, Ar^2 is substituted by 0 to 3 substituents, more preferably 1 or 2 substituents.

Preferably, Ar^2 represents optionally substituted phenyl, isoxazolyl, thiazolyl or thienyl. More preferably, Ar^2 represents optionally substituted phenyl.

Preferably, Ar^2 represents optionally substituted phenyl whereby one or more optional substituents are selected from halogen (such as chloro, e.g. 4-chloro, or fluoro, e.g. 4-fluoro, 2,4-difluoro or 3,4-difluoro), C_{1-6} alkoxy (such as methoxy, e.g. 3-methoxy or 4-methoxy), trifluoromethyl (e.g. 3-trifluoromethyl or 4-trifluoromethyl) and trifluoromethoxy or Ar^2 represents optionally substituted thiazolyl whereby one or more optional substituents are selected from C_{1-6} alkyl (such as methyl, e.g. 2-methyl).


Preferably, R^4 and R^5 independently represent hydrogen or C_{1-4} alkyl. More preferably, R^4 and R^5 independently represent hydrogen or methyl.

Preferably, R^6 and R^7 independently represent hydrogen, fluoro or methyl. More preferably, R^6 and R^7 independently represent hydrogen.

Preferably, R^8 represents hydrogen or methyl. More preferably, R^8 represents hydrogen.

Preferably, p represents 0.


In a first aspect of the invention, there is provided a compound of formula (IA):



or a pharmaceutically acceptable derivative thereof wherein groups A, B, Ar^1 , Ar^2 , Y and R^1 to R^3 have any of the meanings as given hereinbefore. For compounds of formula (IA), R^2 is preferably hydrogen or methoxy.

In a further aspect of the invention, there is provided a compound of the formula (IA) as defined above with the proviso that when m represents 1 and n represents 2 or m represents 2 and n represents 1 and R^2 represents halogen, C_{1-6} alkyl or C_{1-6} alkoxy, Y is other than a bond.

In a further aspect of the invention, Y is a bond and there is provided a compound of formula (IB):




or a pharmaceutically acceptable derivative thereof wherein groups A, B, Ar¹, Ar² and R¹ to R³ have any of the meanings as given hereinbefore.

In a further aspect of the invention, there is provided a compound of the formula (IB) as defined above with the proviso that when m represents 1 and n represents 2 or m represents 2 and n represents 1, R² is other than halogen, C₁₋₆alkyl or C₁₋₆alkoxy.

In a further aspect of the invention, Y is a bond and there is provided a compound of formula (IC):

10




or a pharmaceutically acceptable derivative thereof wherein the groups A, B, Ar² and R¹ to R³ have any of the meanings as given hereinbefore.

In a further aspect of the invention, there is provided a compound of the formula (IC) as defined above with the proviso that when m represents 1 and n represents 2 or m represents 2 and n represents 1, R² is other than halogen, C₁₋₆alkyl or C₁₋₆alkoxy.

In a further aspect of the invention, Y is a bond and there is provided a compound of formula (ID):

20




or a pharmaceutically acceptable derivative thereof wherein groups A, B, R¹ to R³ have any of the meanings as given hereinbefore and the groups R, R' and R'' represent up to three optional substituents on the phenyl ring as defined hereinbefore for the group Ar².

In a further aspect of the invention, there is provided a compound of the formula (ID) as defined above with the proviso that when m represents 1 and n represents 2 or m represents 2 and n represents 1, R² represents halogen, C₁₋₆alkyl or C₁₋₆alkoxy.

30


In a further aspect of the invention, there is provided a compound of formula (IE):

8



or a pharmaceutically acceptable derivative thereof wherein groups Ar¹, Ar², Y and R¹ to R³ have any of the meanings as given hereinbefore.

In a further aspect of the invention, there is provided a compound of formula (IF):




5 or a pharmaceutically acceptable derivative thereof wherein groups Ar¹, Ar², Y and R¹ to R³ have any of the meanings as given hereinbefore.

In a further aspect of the invention, there is provided a compound of the formula (IF) as defined above with the proviso that when R² represents halogen, C₁₋₆alkyl or C₁₋₆alkoxy, Y is other than a bond.

10

In another aspect of the invention, there is provided a compound of formula (IG):




or a pharmaceutically acceptable derivative thereof wherein groups Ar¹, Ar², Y and R¹ to R³ have any of the meanings as given hereinbefore.

15


In a further aspect of the invention, there is provided a compound of the formula (IG) as defined above with the proviso that when R² represents halogen, C₁₋₆alkyl or C₁₋₆alkoxy, Y is other than a bond.

20 In another aspect of the invention, there is provided a compound of formula (IH):



or a pharmaceutically acceptable derivative thereof wherein groups Ar¹, Ar², Y and R¹ to R³ have any of the meanings as given hereinbefore.

In another aspect of the invention, there is provided a compound of formula (IJ):



or a pharmaceutically acceptable derivative thereof wherein groups Ar² and R¹ to R³ have any of the meanings as given hereinbefore.

10 Particular compounds according to the invention include those incorporated in Table 1 and those specifically exemplified and named hereinafter:-


- 5-(4-Chlorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(3-Methoxyphenyl)-thiophene-2-sulfonic acid(8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 15 5-(4-Methoxyphenyl)-thiophene-2-sulfonic acid(8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(3,4-Difluorophenyl)-thiophene-2-sulfonic acid(8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 20 5-(2,4-Difluorophenyl)-thiophene-2-sulfonic acid(8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(3-Chlorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(3-Fluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 25 5-(4-Trifluoromethylphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(3-Trifluoromethylphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 30 5-(4-Fluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(4-Fluorophenyl)-thiophene-2-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 5-(4-Chlorophenyl)-thiophene-2-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;
- 35 5-Isoxazol-3-yl-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-amide;
- 5-(2-Methylthiazol-5-yl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-amide;

- [2,3]Bithiophenyl-5-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)-amide;
 5-(4-Chlorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-yl)amide;
 5 5-(4-Fluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-yl)amide;
 5-(2,4-Difluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-yl)amide; and
 5-(3,4-Difluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1H-3-benzazepin-7-yl)amide.
 10


These compounds may be in the form of their free base or pharmaceutically acceptable salts thereof, particularly the monohydrochloride salt.

- 15 The present invention also provides a general process (A) for preparing compounds of formula (I) which process comprises:

reacting a compound of formula (II)



- 20 with a compound of formula (III)




- 25 wherein Ar¹, Ar² and Y are as hereinbefore defined and R^{1'}-R^{3'} represent R¹ to R³ as hereinbefore defined or are groups that may be readily convertible to R¹ to R³.

- 30 This general method (A) can be conveniently performed by mixing the two components in a solvent such as pyridine or dichloromethane (in the presence of a base), at 0°C.


The present invention also provides a general process (B) for preparing compounds of formula (I) wherein Y is a bond, which process comprises:

reacting a compound of formula (IV)

11



with an aryl boronic acid of formula (V)



5


wherein X is a leaving group, such as iodo, bromo or triflate, Ar¹ and Ar² are as hereinbefore defined and R^{1'}-R^{3'} represent R¹ to R³ as hereinbefore defined or are groups that may be readily convertible to R¹ to R³, under standard Suzuki conditions, e.g. treatment of compound (IV) with 4-chlorobenzenboronic acid in toluene containing aqueous sodium carbonate and a catalytic amount of Pd (PPh₃)₄, at reflux under argon.

10

The present invention also provides a general process (C) for preparing compounds of formula (I) which process comprises:

15

converting a compound of formula (I)



20

wherein A, B, Ar¹, Ar², Y and R¹ to R³ are as hereinbefore defined, into another compound of formula (I) by substituting the group R¹ or the group R³ using conventional techniques.

25

Interconversion of one of the R^{1'} to R^{3'} group to the corresponding R¹ to R³ groups another typically arises when one compound of formula (I) is used as the immediate precursor of another compound of formula (I), or when it is easier to introduce a more complex or reactive substituent at the end of a synthetic sequence.

30

For example, conversion of R^{1'} from a t-butoxycarbonyl (BOC) group to hydrogen is conducted by the treatment of the N-BOC protected compound with hydrogen chloride in ethanol or dioxan at room temperature.

Conversion of R^{1'} from hydrogen to an alkyl group is conducted by the treatment of the sulfonamide NH compound with the appropriate aldehyde in dichloroethane in the presence of a reducing agent, such as sodium triacetoxyborohydride, or by the treatment of the NH compound with the appropriate alkyl halide, such as

iodomethane, under standard alkylation conditions (potassium carbonate in DMF at 60°C).

Conversion of R³ from hydrogen to an alkyl group is conducted by the treatment of the sulfonamide NH compound with the appropriate alcohol, such as methanol, under Mitsunobu conditions i.e. treatment with diisopropyl azodicarboxylate/triphenylphosphine and methanol in tetrahydrofuran at room temperature.

Compounds of formula (II) are known in the literature or may be prepared by known processes, for example, reduction of the corresponding nitro compound as disclosed in WO 99/14197, or by procedures analogous to these procedures. Suitable examples of an R¹ protecting group are trifluoroacetyl or the t-butoxycarbonyl (BOC) group.

Compounds of formula (III) are commercially available or may be prepared by established procedures, for example chlorosulfonylation of a suitable substituted aromatic precursor, using chlorosulfonic acid, for example as described in J. Med. Chem., 2000, 43, 156-166.

Compounds of formula (IV) may be prepared from compounds of formula (II) by the treatment with the appropriate heteroaryl sulfonyl chloride using standard conditions, for example in pyridine or dichloromethane in the presence of a base such as triethylamine at room temperature.

Compounds of formula (V) are commercially available or may be prepared by known methodology, for example lithiation of a suitable substituted bromoheteroaryl at low temperature followed by quenching with tri-isopropylborate and acidic hydrolysis of the reaction product.

Compounds of formula (I) have been found to exhibit affinity for dopamine receptors, in particular the D₃ and D₂ receptors, and are useful in the treatment of disease states which require modulation of such receptors, such as psychotic conditions. Many of the compounds of formula (I) have also been found to have greater affinity for dopamine D₃ than for D₂ receptors. The therapeutic effect of currently available antipsychotic agents (neuroleptics) is generally believed to be exerted via blockade of D₂ receptors; however this mechanism is also thought to be responsible for undesirable extrapyramidal side effects (eps) associated with many neuroleptic agents. Without wishing to be bound by theory, it has been suggested that blockade of the dopamine D₃ receptor may give rise to beneficial antipsychotic activity without significant eps. (see for example Sokoloff et al, Nature, 1990; 347: 146-151; and Schwartz et al, Clinical Neuropharmacology, Vol 16, No. 4, 295-314, 1993). Additionally, certain compounds of formula (I) have antagonist affinity for the

serotonin 5-HT_{2A}, 5-HT_{2C} and 5-HT₆ receptors. These additional properties may give rise to enhanced anti-psychotic activity (e.g. improved effects on cognitive dysfunction) and/or reduced eps.

5 The compounds of formula (I) are of use as antipsychotic agents for example in the treatment of schizophrenia, schizo-affective disorders, psychotic depression, mania, paranoid and delusional disorders. Furthermore, they could have utility as adjunct therapy in Parkinsons Disease, particularly with compounds such as L-DOPA and possibly dopaminergic agonists, to reduce the side effects experienced with these
10 treatments on long term use (e.g. see Schwartz et al., Brain Res. Reviews, 1998, 26, 236-242). From the localisation of D₃ receptors, it could also be envisaged that the compounds could also have utility for the treatment of substance abuse where it has been suggested that D₃ receptors are involved (e.g. see Levant, 1997, Pharmacol. Rev., 49, 231-252). Examples of such substance abuse include alcohol, cocaine,
15 heroin and nicotine abuse. Other conditions which may be treated by the compounds include dyskinetic disorders such as Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias; depression; anxiety, cognitive impairment including memory disorders such as Alzheimers disease, eating disorders, obesity, sexual dysfunction, sleep disorders, emesis, movement disorders, obsessive-compulsive disorders, amnesia, aggression, autism, vertigo, dementia, circadian
20 rhythm disorders and gastric motility disorders e.g. IBS.

The invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable derivative thereof in therapy.

25 The invention also provides a compound of formula (I) or a pharmaceutically acceptable derivative thereof for use in the treatment of a condition in a mammal for which modulation (especially antagonism/inhibition) of dopamine receptors (especially dopamine D₃ receptors) and/or serotonin receptors (especially 5-HT₆, 5-HT_{2A} and 5-HT_{2C}) is beneficial.
30

A preferred use for dopamine/serotonin antagonists according to the present invention is in the treatment of psychoses such as schizophrenia or in the treatment of substance abuse.

35 Therefore, also provided is a compound of formula (I) or a pharmaceutically acceptable derivative thereof for use in the treatment of a psychotic condition (e.g. schizophrenia) or substance abuse in a mammal.

40 The invention also provides the use of a compound of formula (I) or a pharmaceutically acceptable derivative thereof in the manufacture of a medicament for the treatment of a condition in a mammal for which modulation (especially

antagonism/inhibition) of dopamine receptors (especially dopamine D₃ receptors) and/or serotonin receptors (especially 5-HT₆, 5HT_{2A} and 5-HT_{2C}) is beneficial.

5 Also provided is the use of a compound of formula (I) or a pharmaceutically acceptable derivative thereof in the manufacture of a medicament for the treatment of a psychotic condition (e.g. schizophrenia) or substance abuse in a mammal.

10 In a further aspect therefore the present invention provides a method of treating a condition for which modulation (especially antagonism/inhibition) of dopamine receptors (especially dopamine D₃ receptors) and/or serotonin receptors (especially 5-HT₆, 5HT_{2A} and 5-HT_{2C}) is beneficial, which comprises administering to a mammal (e.g. human) in need thereof an effective amount of a compound of formula (I) or a pharmaceutically (i.e. physiologically) acceptable derivative thereof. Such conditions in particular include psychoses/psychotic conditions such as schizophrenia, and
15 substance abuse.

Thus, a still further aspect the invention provides a method of treating a psychotic condition (e.g. schizophrenia) or substance abuse which comprises administering to a mammal (e.g. human) in need thereof an effective amount of a compound of
20 formula (I) as herein defined or a pharmaceutically acceptable derivative thereof.

Also provided is a compound of formula (I) or a pharmaceutically acceptable derivative thereof for use as an active therapeutic substance in a mammal, e.g. for use in the treatment of any of the conditions described herein.
25

"Treatment" includes prophylaxis, where this is appropriate for the relevant condition(s).

For use in medicine, the compounds of the present invention are usually
30 administered as a standard pharmaceutical composition. The present invention therefore provides in a further aspect a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically (i.e. physiologically) acceptable derivative thereof and a pharmaceutically (i.e. physiologically) acceptable carrier. The pharmaceutical composition can be for use in the treatment of any of the
35 conditions described herein.

The compounds of formula (I) may be administered by any convenient method, for example by oral, parenteral (e.g. intravenous), buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted
40 accordingly.

The compounds of formula (I) and their pharmaceutically acceptable derivatives which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

5 A liquid formulation will generally consist of a suspension or solution of the compound or pharmaceutically acceptable derivative in a suitable liquid carrier(s) for example an aqueous solvent such as water, ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

10

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

15 A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion
20 or suspension then filled into a soft gelatin capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or pharmaceutically acceptable derivative in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis
25 oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine
30 suspension of the active substance in a pharmaceutically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted
35 with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

40

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.

- 5 Compositions suitable for transdermal administration include ointments, gels and patches.

Preferably the composition is in unit dose form such as a tablet, capsule or ampoule. Each dosage unit for oral administration contains preferably from 1 to 250 mg (and
10 for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I) or a pharmaceutically acceptable derivative thereof calculated as the free base.

The pharmaceutically acceptable compounds of the invention will normally be
15 administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 500 mg, preferably between 10 mg and 400 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1
20 and 25 mg of the compound of the formula (I) or a pharmaceutically acceptable derivative thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

Biological Test Methods

25 Binding experiments on cloned dopamine (e.g. D2 and D3) receptors

The ability of the compounds to bind selectively to human D2/D3 dopamine receptors can be demonstrated by measuring their binding to cloned receptors. The inhibition constants (K_i) of test compounds for displacement of [125 I]-iodosulpride binding to
30 human D2/D3 receptors expressed in CHO cells were determined as follows. The cell lines were shown to be free from bacterial, fungal and mycoplasmal contaminants, and stocks of each were stored frozen in liquid nitrogen. Cultures were grown as monolayers or in suspension in standard cell culture media. Cells were recovered by scraping (from monolayers) or by centrifugation (from suspension cultures), and were washed two or three times by suspension in phosphate buffered
35 saline followed by collection by centrifugation. Cell pellets were stored frozen at -80°C. Crude cell membranes were prepared by homogenisation followed by high-speed centrifugation, and characterisation of cloned receptors achieved by radioligand binding.

40 Preparation of CHO cell membranes: Cell pellets were gently thawed at room temperature, and resuspended in about 20 volumes of ice-cold Extraction buffer; 5mM EDTA, 50mM Trizma pre-set crystals (pH7.4@37°C), 1mM $MgCl_2$, 5mM KCl and 120mM NaCl. The suspension was homogenised using an Ultra-Turrax at full

speed for 15 seconds. The homogenate was centrifuged at 18,000 r.p.m for 15 min at 4°C in a Sorvall RC5C centrifuge. Supernatant was discarded, and homogenate re-suspended in extraction buffer then centrifugation was repeated. The final pellet was resuspended in 50mM Trizma pre-set crystals (pH 7.4 @ 37°C) and stored in 1ml aliquot tubes at -80°C (D2 = 3.0E+08 cells, D3 = 7.0E+07 cells and D4 = 1.0E+08 cells). The protein content was determined using a BCA protocol and bovine serum albumin as a standard (Smith, P. K., et al., Measurement of protein using bicinchoninic acid. Anal. Biochem. 150, 76-85 (1985)).

Binding experiments: Crude D2/D3 cell membranes were incubated with 0.03nM [¹²⁵I]-Iodosulpride (~2000 Ci/mmol; Amersham, U. K., and the test compound in a buffer containing 50mM Trizma pre-set crystals (pH 7.4 @ 37°C), 120mM NaCl, 5mM KCl, 2mM CaCl₂, 1mM MgCl₂, 0.3% (w/v) bovine serum albumin. The total volume is 0.2ml and incubated in a water bath at 37°C for 40 minutes. Following incubation, samples were filtered onto GF/B Unifilters using a Canberra Packard Filtermate, and washed four times with ice-cold 50mM Trizma pre-set crystals (pH 7.4 @ 37°C). The radioactivity on the filters was measured using a Canberra Packard Topcount Scintillation counter. Non-specific binding was defined with 10µM SKF-102161 (YM-09151). For competition curves, 10 serial log concentrations of competing cold drug were used (Dilution range: 10µM-10pM). Competition curves were analysed using Inflexion, an iterative curve fitting programme in Excel. Results were expressed as pK_i values where
$$pK_i = -\log_{10}[K_i]$$

The exemplified compounds have pK_i values within the range of 7.4-8.9 at the dopamine D₃ receptor.
The exemplified compounds have pK_i values within the range of 6.4-7.8 at the dopamine D₂ receptor.

Binding experiments on cloned 5-HT₆ receptors

Compounds can be tested following the procedures outlined in WO 98/27081.

The exemplified compounds have pK_i values within the range of 7.5-8.7 at the serotonin 5-HT₆ receptor.

Binding experiments on cloned 5-HT_{2C} receptors

Compounds can be tested following the procedures outlined in WO 94/04533 and *British Journal of Pharmacology* (1996) 117, 427-434.

The exemplified compounds have pK_i values within the range of 5.7-7.5 at the serotonin 5-HT_{2C} receptor.

Binding experiments on cloned 5-HT_{2A} receptors

Compounds can be tested following the procedures outlined in *British Journal of Pharmacology* (1996) 117, 427-434.


The exemplified compounds have pK_i values within the range of 6.0-7.8 at the serotonin 5-HT_{2A} receptor.

The invention is further illustrated by the following non-limiting examples:

5

Description 1

7-Amino-8-methoxy-1,2,4,5-tetrahydro-3-benzazepine-3-carboxylic acid tert-butyl ester (D1)



10 7-Hydroxy-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid tert-butyl ester (D1a)

7-Methoxy-2,3,4,5-tetrahydro-1H-3-benzazepine (10 g) in 48% aqueous hydrobromic acid (350 ml) was allowed to stir at 100°C for 4 h. The mixture was allowed to cool to 20°C then evaporated to dryness, giving the crude hydroxy compound as a brown solid (14.5 g). This solid was dissolved in tetrahydrofuran (100 ml) and water (70 ml) and triethylamine (8 g) was added dropwise, followed by a solution of di-tert-butyl dicarbonate (14 g) in tetrahydrofuran (20 ml). The resulting mixture was allowed to stir at 20°C for 16 h then partitioned between ethyl acetate (200 ml) and water (200 ml). The aqueous layer was extracted with ethyl acetate (100 ml). The combined organic extracts were washed with saturated aqueous sodium bicarbonate (100 ml), dried over anhydrous sodium sulfate and evaporated to dryness. The resulting oil was purified by chromatography over silica gel, eluting with 10-30% ethyl acetate in hexane, affording the title compound D1a as a white solid (8 g), MS (API⁺): Found 164 (MH⁺-Boc). C₁₅H₂₁NO₃ requires 263. ¹H NMR: δ CDCl₃ 1.48 (9H, s), 2.75-2.87 (4H, m), 3.40-3.60 (4H, m), 4.95 (1H, s), 6.50-6.62 (2H, m), 6.96 (1H, d).

20 7-Methoxy-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid tert-butyl ester (D1b)

Reaction of the phenol D1a with potassium carbonate/methyl iodide in dimethylformamide afforded the title compound. MH⁺ 278.

30 7-Methoxy-8-nitro-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid tert-butyl ester (D1c)


Nitration of D1b was carried out using a solution of nitric acid and acetic anhydride; the crude product was purified by chromatography on silica gel using EtOAc/n-hexane as eluant to afford the title compound D1c. M⁺ - C(CH₃)₃ + 2H = 267

35 7-Amino-8-methoxy-1,2,4,5-tetrahydro-3-benzazepine-3-carboxylic acid tert-butyl ester (D1)

Hydrogenation of D1c at 50 psi in ethanol over 10% palladium on charcoal at room temperature afforded the title compound D1. MH^+ 293.

Description 2

- 5 **7-Amino-1,2,4,5-tetrahydro-3-benzazepine-3-carboxylic acid *tert*-butyl ester (D2)**



This compound was prepared using the procedure described in EP 284384 i.e.

- 10 **7-Nitro-1,2,4,5-tetrahydro-3H-3-benzazepine (D2a)**

1,2,4,5-Tetrahydro-3H-benzazepine (1 g) (See P. Ruggli et al., *Helv. Chim. Acta*, 18, 1388, [1935]) was added slowly dropwise to stirred fuming nitric acid (25 ml) at -10°C. Stirring was continued at -10°C for 1 hour and the reaction mixture was then poured onto ice, the precipitate collected by filtration and dried to give the title compound as the nitrate salt, 1.4g. This was suspended in water, cooled to 5°C and neutralised with 5M sodium hydroxide. The precipitate was collected by filtration, recrystallised from water and dried, affording the title compound D1a as a white solid (0.6 g)

- 20 **3-Tert-butoxycarbonyl-7-nitro-1,2,4,5-tetrahydro-3H-3-benzazepine (D2b)**

A solution of di-*t*-butyldicarbonate (2.18 g) in dry dichloromethane (15 ml) was added dropwise to a stirred solution of 7-nitro-1,2,4,5-tetrahydro-3H-3-benzazepine (1.92 g) in dry dichloromethane (40 ml) at 0°C. After 18 h at room temperature the solvent was evaporated, giving an oil. This oil was dissolved in dichloromethane, washed twice with saturated aqueous sodium bicarbonate, three times with 1M hydrochloric acid and twice with brine. The organic solution was dried and evaporated giving an oil D2b, 2.33 g.


- 30 **7-Amino-1,2,4,5-tetrahydro-3-benzazepine-3-carboxylic acid *tert*-butyl ester (D2)**

A solution of 3-*tert*-butoxycarbonyl-7-nitro-1,2,4,5-tetrahydro-3H-3-benzazepine (2.1 g) was stirred under a hydrogen atmosphere (50 p.s.i.) in ethanol (40 ml) containing 5% Pd/C (0.21 g) for 3 hours. The catalyst was removed by filtration and the solvent evaporated to give the title compound D2 as a low-melting solid, 2.0 g. MH^+ 263

35

Description 3


Amino-dimethylamino-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert*-butyl ester (D3)



This compound was prepared according to the procedure described in WO 03/068752A1

5 Description 4


7-(5-Bromo-thiophene-2-sulfonylamino)-8-methoxy-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert* -butyl ester (D4)



To an ice bath cooled solution of the methoxy aniline intermediate D1 (2.0 g, 0.0068 moles) in dry pyridine was added dropwise a solution of 5-bromothiophene-2-sulfonyl chloride (2.01 g, 0.0077 moles) in dichloromethane. The mixture was stirred for 1 hour at room temperature. The mixture was evaporated to dryness then purified by chromatography over silica, eluting with hexane up to 50% EtOAc/hexane, affording the title compound D4 as a white foam (3.18 g, 90%). M-H 517. ¹H NMR: δ CDCl₃ 1.48 (9H, s), 2.81 (4H, m), 3.49 (4H, m), 3.67 (3H, s), 6.60 (1H, s), 6.90 (1H, s), 6.94 (1H, d), 7.19 (1H, d), 7.30 (1H, s).

Description 5

7-(5-Bromo-thiophene-2-sulfonylamino)-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert*-butyl ester (D5)




This compound was prepared using a similar procedure to that described for D4 above. MH⁺ 488.

25 Description 6

(5-Bromo-thiophene-2-sulfonylamino)-dimethylamino-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert* -butyl ester (D6)


21



To an ice bath cooled solution of the intermediate D3 (2.0 g, 0.0065 moles) in dry pyridine was added dropwise a solution of 5-bromothiophene-2-sulfonyl chloride (1.88 g, 0.0072 moles) in dichloromethane. The mixture was stirred for 1 hour at room temperature. The mixture was evaporated to dryness then purified by chromatography over silica, eluting with hexane up to 50% EtOAc/hexane, affording the title compound D6 as a brown solid (2.1 g, 90%). M-H 532. ¹H NMR: δ MeOD 1.45 (9H,s), 2.76 (2H, m), 3.02 (2H, m), 3.30 (6H,s), 3.48 (2H, m), 3.58 (2H, m), 6.47 (1H, s), 7.26 (2H, m), 7.76 (1H, s).

Description 7

7-[5-(4-Chloro-phenyl)-thiophene-2-sulfonylamino]-8-methoxy-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert*-butyl ester (D7)




A solution of the bromo intermediate D5 (1.20 g, 0.0023 moles), 4-chlorophenylboronic acid (0.55 g, 0.0034 moles), K₂CO₃ (2M solution, 15 ml), ethanol (15 ml) and toluene (50 ml) at room temperature was degassed by bubbling argon through the solution for 10 minutes. Pd(PPh₃)₄ (0.4 g, 0.000375 moles) was added and the mixture heated at 60°C under argon for 4 hours. Upon cooling the mixture was partitioned between water and ethyl acetate. The aqueous phase was re-extracted with ethyl acetate (x3) and the combined organic phase washed with water, brine then dried over anhydrous MgSO₄. The solution was evaporated to dryness then purified by chromatography over silica eluting, with hexane up to 50% EtOAc/hexane, affording the title compound D7 as a white foam (1.09 g, 86%). M-H 547. ¹H NMR: δ CDCl₃ 1.47 (9H, s) 2.83(4H, m), 3.50 (4H, m), 3.66 (3H,s), 6.55 (1H, s), 6.96 (1H, s), 7.12 (1H, d), 7.33-7.50 (5H, m).

Description 8

[5-(4-Chloro-phenyl)-thiophene-2-sulfonylamino]-dimethylamino-1,2,4,5-tetrahydro-benzo[d]azepine-3-carboxylic acid *tert*-butyl ester (D8)


22



A solution of the bromo intermediate D6 (0.2 g, 0.00037 moles), 4-chlorophenylboronic acid (0.070 g, 0.00045 moles), K_2CO_3 (2M solution, 1 ml), ethanol (1 ml) and toluene (2 ml) at room temperature was degassed by bubbling argon through the solution for 10 minutes. $Pd(PPh_3)_4$ (0.1 g, 0.000086 moles) was added and the mixture heated at 60°C under argon for 24 hours. Upon cooling the mixture was partitioned between water and ethyl acetate. The aqueous phase was re-extracted with ethyl acetate (x3) and the combined organic phase washed with water, brine then dried over anhydrous $MgSO_4$. The solution was evaporated to dryness then purified by chromatography over silica eluting, with hexane up to 50% EtOAc/hexane, affording the title compound D8 as a off white foam, which was used directly in the next step. M-H 563.

15 Description 9


5-(4-Chloro-phenyl)-thiophene-2-sulfonic acid (8-methoxy-2,3, 4,5-tetrahydro-1-*H*-benzo[d]azepin-7-yl)-amide (D9)



A solution of the Boc-protected amine D7 (1.05g, 0.00199 moles) in 1,4-dioxane (20 ml) and 4M HCl in dioxane (10 ml) was stirred under argon at room temperature overnight. The resultant mixture was evaporated to dryness to give the hydrochloride salt of the title compound D9 as a cream solid (0.96g, 100%). M-H 448.5. 1H NMR: δ DMSO- d_6 3.08 (4H, m), 3.32 (4H, br s), 3.52 (3H, s), 6.85 (1H, s), 7.10 (1H, s), 7.41 (1H, d), 7.49-7.53 (3H, m), 7.69 (1H, s), 7.73 (1H, s), 9.27 (2H, br s), 9.74 (1H, br s).

Description 10


5-(4-Chloro-phenyl)-thiophene-2-sulfonic acid (8-dimethylamino-2,3, 4,5-tetrahydro-1-*H*-benzo[d]azepin-7-yl)-amide (D10)



A solution of the intermediate D8 in methanol (1 ml) and 4M HCl in dioxane (5 ml) was stirred under argon at room temperature overnight. The resultant mixture was evaporated to dryness to give the hydrochloride salt of the title compound D10 as a off white foam, which was used directly in the next step. M-H 463.

Example 1

5-(4-Chloro-phenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1-*H*-benzo[*d*]azepin-7-yl)-amide (E1)




To a suspension of the amine hydrochloride salt D9 (0.96g, 0.00198 moles) in 1,2-dichloroethane (50 ml) at room temperature was added triethylamine (5 ml, excess) followed by 37% aqueous formaldehyde (5 ml, excess). After vigorous stirring for 5 minutes, sodium triacetoxyborohydride (5g, excess) was added portionwise over 5 minutes and the resultant solution stirred for a further 2 hours. The reaction was partitioned between water and dichloromethane and the organic phase washed with water, brine and dried over anhydrous MgSO_4 . The solution was evaporated to dryness affording the title compound E1 as a pale yellow solid (0.993g, 100%). M-H 461. ^1H NMR: δ CDCl_3 2.67-2.88 (6H, m), 3.60-3.95 (8H, m), 6.58 (1H, s), 7.07 (1H, s), 7.13 (1H, d), 7.33-7.50 (6H, m).

Examples 2-12 were prepared using analogous procedures to Example 1 using the appropriate starting materials, with the products being isolated as either free bases or hydrochloride salts. Examples 13 to 15 were prepared using analogous procedures to those described in Descriptions 3, 4, 6 and Example 1 from the commercially available sulfonyl chlorides with the products being isolated as either free bases or hydrochloride salts. All ^1H NMR are consistent with the structures shown.

Example 16

5-(4-Chloro-phenyl)-thiophene-2-sulfonic acid (dimethylamino-methyl-2,3,4,5-tetrahydro-1-*H*-benzo[*d*]azepin-7-yl)-amide (E16)



To a suspension of the intermediate D10, in 1,2-dichloroethane (1 ml) at room temperature was added triethylamine (0.3, excess) followed by 37% aqueous formaldehyde (0.2, excess). After vigorous stirring for 5 minutes, sodium triacetoxymethylborohydride (100 mg, excess) was added portionwise over 5 minutes and the resultant solution stirred for a further 2 hours. The reaction was partitioned between water and dichloromethane and the organic phase washed with water, brine and dried over anhydrous MgSO_4 . The solution was evaporated to dryness affording the title compound E16 as a pale yellow solid (20 mg, 99%). M-H 477. ^1H NMR: δ CDCl_3 2.36 (3H, s), 2.44 (6H, s), 2.54 (4H,m), 2.84 (2H, m), 2.91 (2H, m), 6.87 (1H, s), 7.12 (1H, d), 7.33-7.51 (6H, m).

Examples 17-19 were prepared using analogous procedures to Example 16 using the appropriate starting materials, with the products being isolated as either free bases or hydrochloride salts. All ^1H NMR are consistent with the structures shown.

All of the compounds listed below in Table 1 relate to compounds of formula (IJ):





Table 1

Example	R ¹	R ²	R ³	Ar ⁴	MH ⁺
1	Me	MeO	H	4-chlorophenyl	461
2	Me	MeO	H	3-methoxyphenyl	459
3	Me	MeO	H	4-methoxyphenyl	459
4	Me	MeO	H	3,4-difluorophenyl	465
5	Me	MeO	H	2,4-difluorophenyl	465
6	Me	MeO	H	3-chlorophenyl	463
7	Me	MeO	H	3-fluorophenyl	447
8	Me	MeO	H	4-trifluoromethylphenyl	497
9	Me	MeO	H	3-trifluoromethylphenyl	497
10	Me	MeO	H	4-fluorophenyl	447
11	Me	H	H	4-fluorophenyl	417
12	Me	H	H	4-chlorophenyl	433
13	Me	MeO	H	3-isoxazolyl	404
14	Me	MeO	H	2-methyl-thiazol-5-yl	450
15	Me	H	H	3-thienyl	405
16	Me	NMe ₂	H	4-chlorophenyl	477
17	Me	NMe ₂	H	4-fluorophenyl	460
18	Me	NMe ₂	H	2,4-difluorophenyl	478
19	Me	NMe ₂	H	3,4-difluorophenyl	478

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

CLAIMS

1. A compound of formula (I):



wherein

A and B represent the groups $-(CH_2)_m-$ and $-(CH_2)_n-$ respectively;

R^1 represents C_{1-6} alkyl;

R^2 represents hydrogen, halogen, hydroxy, cyano, nitro, hydroxy C_{1-6} alkyl, trifluoromethyl, trifluoromethoxy, C_{1-6} alkyl, C_{1-6} alkoxy, $-(CH_2)_pC_{3-6}$ cycloalkyl, $-(CH_2)_pOC_{3-6}$ cycloalkyl, $-COC_{1-6}$ alkyl, $-SO_2C_{1-6}$ alkyl, $-SOC_{1-6}$ alkyl, $-SC_{1-6}$ alkyl, $-CO_2C_{1-6}$ alkyl, $-CO_2NR^4R^5$, $-SO_2NR^4R^5$, $-(CH_2)_pNR^4R^5$, $-(CH_2)_pNR^4COR^5$, an optionally substituted aryl group, an optionally substituted heteroaryl group or an optionally substituted heterocyclyl group;

R^3 represents hydrogen or C_{1-6} alkyl;

Ar^1 represents an optionally substituted heteroaryl group;

Ar^2 represents an optionally substituted phenyl or an optionally substituted heteroaryl group;

Y represents a bond, $-O-$, $-C_{1-6}$ alkyl-, $-CR^6R^7X-$, $-XCR^6R^7-$, $-NR^8CO-$ or $-CONR^8-$;

X represents oxygen, sulfur, $-SO-$ or $-SO_2-$;

R^4 and R^5 each independently represent hydrogen or C_{1-6} alkyl or, together with the nitrogen or other atoms to which they are attached, form an azacycloalkyl ring or an oxo-substituted azacycloalkyl ring;

R^6 and R^7 each independently represent hydrogen, C_{1-6} alkyl or fluoro;

R^8 represents hydrogen or C_{1-6} alkyl;

m and n independently represent an integer selected from 1 and 2;

p independently represents an integer selected from 0, 1, 2 and 3;

or a pharmaceutically acceptable salt, solvate or pharmaceutically acceptable derivative thereof.

2. A compound of formula (I) which is

5-(4-Chlorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;

5-(3-Methoxyphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;

5-(4-Methoxyphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;

5-(3,4-Difluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1H-benzo[d]azepin-7-yl)amide;

- 5-(2,4-Difluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5-(3-Chlorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5 5-(3-Fluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5-(4-Trifluoromethylphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5-(3-Trifluoromethylphenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
10 5-(4-Fluorophenyl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5-(4-Fluorophenyl)-thiophene-2-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
15 5-(4-Chlorophenyl)-thiophene-2-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)amide;
5-(4-Chloro-2-methylphenyl)-thiophene-2-sulfonic acid (2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)-amide;
5-Isoxazol-3-yl-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)-amide;
20 5-(2-Methylthiazol-5-yl)-thiophene-2-sulfonic acid (8-methoxy-3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)-amide;
[2,3']Bithiophenyl-5-sulfonic acid (2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)-amide;
[2,3']Bithiophenyl-5-sulfonic acid (3-methyl-2,3,4,5-tetrahydro-1*H*-benzo[d]azepin-7-yl)-amide;
25 5-(4-Chlorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepin-7-yl)amide;
5-(4-Fluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepin-7-yl)amide;
30 5-(2,4-Difluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepin-7-yl)amide; and
5-(3,4-Difluorophenyl)thiophene-2-sulfonic acid (8-dimethylamino-3-methyl-2,3,4,5-tetrahydro-1*H*-3-benzazepin-7-yl)amide.
- 35 3. A pharmaceutical composition comprising a compound of formula (I) as claimed in claim 1 or claim 2 or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier therefor.
- 40 4. Use of a compound of formula (I) according to claim 1 or claim 2 or a pharmaceutically acceptable derivative thereof in therapy.
5. Use of a compound of formula (I) according to claim 1 or claim 2 for the treatment of a condition which requires modulation of a dopamine receptor.

6. Use of a compound of formula (I) according to claim 5 wherein the condition is schizophrenia or substance abuse.
- 5 7. Use of a compound of formula (I) according to claim 1 or claim 2 in the manufacture of a medicament for the treatment of a condition which requires modulation of a dopamine receptor.
- 10 8. Use of a compound of formula (I) according to claim 7 wherein the condition is schizophrenia or substance abuse.
- 15 9. A method of treating a condition which requires modulation of dopamine receptors which comprises administering to a mammal in need thereof an effective amount of a compound of formula (I) according to claim 1 or claim 2.
10. A method of treating a condition according to claim 9 wherein the condition is schizophrenia or substance abuse.

INTERNATIONAL SEARCH REPORT

Internat

Application No

PCT/EP 03/11174

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/12 C07D413/14 C07D417/14 C07D409/14 A61K31/55
A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 100 53 799 A (BAYER AG) 8 May 2002 (2002-05-08) the whole document	1-10
Y	WO 01 32646 A (SMITHKLINE BEECHAM) 10 May 2001 (2001-05-10) the whole document	1-10
Y	WO 01 45694 A (SMITHKLINE BEECHAM) 28 June 2001 (2001-06-28) the whole document	1-10
Y	EP 0 007 070 A (SMITHKLINE CORP.) 23 January 1980 (1980-01-23) the whole document	1-10
	--- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

30 January 2004

Date of mailing of the international search report

12/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

Intern: Application No
PCT/EP 03/11174

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	WO 03 068752 A (GLAXO GROUP LTD.) 21 August 2003 (2003-08-21) the whole document -----	1-10

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/11174

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 9 and 10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal

Application No

PCT/EP 03/11174

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 10053799	A	08-05-2002	DE 10053799 A1	08-05-2002
WO 0132646	A	10-05-2001	AU 1278701 A	14-05-2001
			WO 0132646 A2	10-05-2001
			EP 1228066 A2	07-08-2002
			JP 2003513085 T	08-04-2003
WO 0145694	A	28-06-2001	AU 2441801 A	03-07-2001
			CA 2394603 A1	28-06-2001
			EP 1248607 A1	16-10-2002
			JP 2003518057 T	03-06-2003
			WO 0145694 A1	28-06-2001
EP 7070	A	23-01-1980	AR 222828 A1	30-06-1981
			AT 2267 T	15-02-1983
			AT 19069 T	15-04-1986
			AU 525789 B2	02-12-1982
			AU 4861379 A	07-02-1980
			CA 1115271 A1	29-12-1981
			CS 213381 B2	09-04-1982
			DD 147355 A5	01-04-1981
			DE 2964530 D1	24-02-1983
			DE 2967590 D1	15-05-1986
			DK 278379 A ,B,	08-01-1980
			EP 0007070 A1	23-01-1980
			EP 0080012 A1	01-06-1983
			ES 482276 A1	01-04-1980
			ES 482281 A1	01-04-1980
			ES 482282 A1	01-04-1980
			FI 792125 A ,B,	08-01-1980
			GR 69675 A1	07-07-1982
			HU 179315 B	28-09-1982
			IE 48655 B1	03-04-1985
			IE 48654 B1	03-04-1985
			IL 57532 A	31-08-1982
			IT 1193190 B	02-06-1988
			JP 1053271 B	13-11-1989
			JP 55011584 A	26-01-1980
			MY 93387 A	31-12-1987
			NO 792267 A ,B,	08-01-1980
			NZ 190745 A	23-01-1981
			PH 15754 A	18-03-1983
			PL 216908 A1	11-08-1980
			PT 69855 A	01-08-1979
			SU 1029827 A3	15-07-1983
			US 4265890 A	05-05-1981
			ZA 7902785 A	27-08-1980
WO 03068752	A	21-08-2003	WO 03068752 A1	21-08-2003
			WO 03068751 A1	21-08-2003